Complements of Multivalued Functions

Stephen Fenner [*]
University of Southern Maine

Frederic Green [†]
Clark University

Steven Homer [‡]
Boston University

Alan L. Selman [§]
State University of New York at Buffalo

Thomas Thierauf [¶]
Universität Ulm

Heribert Vollmer [‖]
Universität Würzburg

Abstract

We study the class coNPMV *of complements of* NPMV *functions. Though defined symmetrically to* NPMV *this class exhibits very different properties. We clarify the complexity of* coNPMV *by showing that it is essentially the same as that of* $\mathrm{NPMV}^{\mathrm{NP}}$. *Complete functions for* coNPMV *are exhibited and central complexity-theoretic properties of this class are studied. We show that computing maximum satisfying assignments can be done in* coNPMV, *which leads us to a comparison of* NPMV *and* coNPMV *with Krentel's classes* Max P *and* Min P. *The difference hierarchy for* NPMV *is related to the query hierarchy for* coNPMV. *Finally, we examine a functional analogue of Chang and Kadin's relationship between a collapse of the Boolean hierarchy over* NP *and a collapse of the polynomial time hierarchy.*

1. Introduction

Consider the complexity class NPMV of partial multivalued functions that are computed nondeterministically in polynomial time. As this class captures the complexity of computing witnesses of sets in NP, by studying this class, and more generally, by studying relations between such complexity classes of partial multivalued functions, we directly contribute to understanding the complexity of computing witnesses. It is well-known that a partial multivalued function f belongs to NPMV if and only if it is polynomial length-bounded and $\mathrm{graph}(f) = \{\langle x, y\rangle \mid y \text{ is a value of } f(x)\}$ belongs to NP.

Now consider the class coNPMV. We will give a formal definition in the preliminaries section below. For now, let us state that a partial multivalued function f belongs to coNPMV if and only if it is polynomial length-bounded and $\mathrm{graph}(f)$ belongs to coNP. Given this symmetry, graphs of functions in NPMV are in NP while graphs of functions in coNPMV are in coNP, and given what we know about NP and coNP, one might expect that coNPMV has essentially the same complexity as NPMV. Indeed, it is easy to see that coNPMV = NPMV if and only if NP = coNP. However, the point of this paper is to show that in many ways coNPMV is a more powerful class than is NPMV. One can derive more information from computing the complement of a function in NPMV than from computing the function. For one example of this phenomena, we prove here that coNPMV is not included in $\mathrm{FP}^{\mathrm{NPMV}}$ unless the polynomial hierarchy collapses. (This is an extension of a result of Fenner *et al.* [FHOS93].) Thus, a coNPMV oracle provides more information than an NPMV oracle. This is surprising, for function oracles, just as set oracles, provide knowledge about both their domains and their co-domains.

We will define many-one reductions between multivalued functions. This will be a straightforward adaptation of the many-one metric reducibility of Krentel [Kre88]. In Section 3, we will consider many-one complete functions for coNPMV.

In Section 4, Consider the partial multivalued function *sat*, defined so that y is a value of $sat(\varphi)$ if and only if y is a satisfying assignment of Boolean formula φ. The function *sat* is complete for NPMV. Nevertheless, in Section 4 we will see that *sat* and similar such functions belong to coNPMV. Even the seemingly more powerful $\mathrm{FP}^{\mathrm{NP}}$-complete function *maxsat*, that gives the maximum satisfying assignment of a formula, is contained in coNPMV. However, we will see

[*]Computer Science Department, University of Southern Maine, Portland, ME 04103. E-mail: fenner@cs.usm.maine.edu. Supported in part by the NSF under grants CCR 92-09833 and CCR 95-01794.

[†]Department of Mathematics and Computer Science, Clark University, Worcester, MA 01610. E-mail: fgreen@black.clarku.edu.

[‡]Computer Science Department, Boston University, Boston, MA 02215. E-mail: homer@cs.bu.edu. Supported in part by the NSF under grant NSF-CCR-9400229.

[§]Department of Computer Science, 226 Bell Hall, State University of New York at Buffalo, Buffalo, NY 14260. E-mail: selman@cs.buffalo.edu. Supported in part by the NSF under grant NSF-CCR-9400229.

[¶]Abteilung Theoretische Informatik, Universität Ulm, Oberer Eselsberg, 89069 Ulm, Germany. Email: thierauf@informatik.uni-ulm.de. Supported in part by DAAD, Acciones Integradas 1995, 322-AI-e-dr.

[‖]Theoretische Informatik, Universität Würzburg, Am Exerzierplatz 3, D-97072 Würzburg, Germany. E-mail: vollmer@informatik.uni-wuerzburg.de. Work supported by the Alexander von Humboldt foundation under a Feodor Lynen scholarship.

that NPMV or $\mathrm{FP}^{\mathrm{NP}}$ are *not* contained in coNPMV, and hence coNPMV is not closed under metric many-one reductions, unless the polynomial time hierarchy collapses. Clearly, these function classes have strange closure properties, which we describe below.

As an upper bound on the complexity of coNPMV, we show that, for any $k \geq 2$,

$$\mathrm{coNPMV} \subseteq \mathrm{NPMV}(2) \subseteq \mathrm{NPMV}(k) \subseteq$$

$$\mathrm{NPMV}(k+1) \subseteq \mathrm{NPMV}(n^{O(1)}) \subseteq \mathrm{NPMV}^{\mathrm{NP}},$$

where $\mathrm{NPMV}(k)$ is the k-th level of the difference hierarchy for NPMV as defined by Fenner *et al.* [FHOS93].

On the other hand, even though there is an infinite hierarchy of complexity classes between coNPMV and $\mathrm{NPMV}^{\mathrm{NP}}$ (the difference hierarchy over NPMV does not collapse unless the polynomial time hierarchy collapses [FHOS93]), our results suggest that the complexity of coNPMV is essentially the same as the complexity of $\mathrm{NPMV}^{\mathrm{NP}}$: we prove in Section 5 that $\mathrm{NPMV}^{\mathrm{NP}} = \pi_2^1 \circ \mathrm{coNPMV}$ (where π_2^1 is the projection function that maps a pair of strings to its first component). It follows that $\mathrm{NPMV}^{\mathrm{NP}}$ is the closure of coNPMV under metric many-one reductions.

In Section 6, we show that if the difference hierarchy for NPMV collapses, then the NPMV oracle hierarchy collapses. This is the functional analogue of the well-known result by Chang and Kadin relating a collapse of the Boolean hierarchy over NP to a collapse of the polynomial time hierarchy.

2. Preliminaries

We fix Σ to be the finite alphabet $\{0,1\}$. Let $<$ denote the standard lexicographic order on Σ^*. By $\langle \cdot, \cdot \rangle$ we denote a standard pairing function on $\Sigma^* \times \Sigma^*$.

We use the standard complexity classes P and NP for (nondeterministic) polynomial time, Σ_k^p and $\Delta_k^p = \mathrm{P}^{\Sigma_{k-1}^p}$ for the levels of the polynomial time hierarchy, and $\mathrm{NP}(k)$ for the levels of the Boolean hierarchy, for $k \geq 1$.

Let f be a relation on $\Sigma^* \times \Sigma^*$. We will call f a *(partial) multivalued function* from Σ^* to Σ^*. By $f(x) \mapsto y$ we denote that $(x,y) \in f$ and say that f *maps x to y*. By set-$f(x)$ we denote the set of outcomes of f on x, set-$f(x) = \{ y \mid f(x) \mapsto y \}$. The *graph of* f is $\mathrm{graph}(f) = \{ \langle x, y \rangle \mid f(x) \mapsto y \}$. The *domain of* f, $\mathrm{dom}(f)$, is the set of x where set-$f(x)$ is nonempty. We will say that f is undefined at x if $x \notin \mathrm{dom}(f)$. The domain of a class $\mathcal{F}$ of functions is $\mathrm{dom}(\mathcal{F}) = \bigcup_{f \in \mathcal{F}} \mathrm{dom}(f)$.

Given partial multivalued functions f and g, define g to be a *refinement of* f if $\mathrm{dom}(g) = \mathrm{dom}(f)$ and $\mathrm{graph}(g) \subseteq \mathrm{graph}(f)$. Let $\mathcal{F}$ and $\mathcal{G}$ be classes of partial multivalued functions. Purely as a convention, if f is a partial multivalued function, we define $f \in_c \mathcal{G}$ if $\mathcal{G}$ contains a refinement of f, and we define $\mathcal{F} \subseteq_c \mathcal{G}$ if for every $f \in \mathcal{F}$, $f \in_c \mathcal{G}$. This notation is consistent with our intuition that $\mathcal{F} \subseteq_c \mathcal{G}$ should entail that the complexity $\mathcal{F}$ is not greater than the complexity of $\mathcal{G}$.

A transducer T is a nondeterministic Turing machine with a read-only input tape, a write-only output tape, read-write work tapes, and accepting states in the usual manner. T computes a value y on an input string x if there is an accepting computation of T on x for which y is the final content of T's output tape. (In this case, we will write $T(x) \mapsto y$.) Such transducers compute partial, multivalued functions. (As transducers do not typically accept all input strings, when we write "function", "partial function" is always intended. If a function f is total, it will always be explicitly noted.)

The following classes of partial functions were first defined by Book, Long, and Selman [BLS84].

- NPMV is the set of all partial, multivalued functions computed by nondeterministic polynomial time-bounded transducers;
- NPSV is the set of all $f \in \mathrm{NPMV}$ that are single-valued;
- FP is the set of all partial functions computed by deterministic polynomial time-bounded transducers.

A function f belongs to NPMV if and only if it is polynomially length-bounded and $\mathrm{graph}(f)$ belongs to NP. The domain of every function in NPMV belongs to NP. An example is *sat*, which maps Boolean formulas to their satisfying assignments.

Fenner *et al.* [FHOS93] define the *difference hierarchy over* NPMV as follows. Let $\mathcal{F}$ be a class of partial multivalued functions. A partial multivalued function f is in co$\mathcal{F}$ if there exist $g \in \mathcal{F}$ and a polynomial p such that for every x,

$$\text{set-}f(x) \;=\; \Sigma^{p(|x|)} - \text{set-}g(x).$$

Let $\mathcal{F}$ and $\mathcal{G}$ be two classes of partial multivalued functions. A partial multivalued function h is in $\mathcal{F} \wedge \mathcal{G}$, respectively $\mathcal{F} \vee \mathcal{G}$, if there exist partial multivalued functions $f \in \mathcal{F}$ and $g \in \mathcal{G}$ such that for every x,

$$\begin{aligned} \text{set-}h(x) &= \text{set-}f(x) \cap \text{set-}g(x), \text{ respectively} \\ \text{set-}h(x) &= \text{set-}f(x) \cup \text{set-}g(x). \end{aligned}$$

Let $\mathcal{F} - \mathcal{G}$ denote $\mathcal{F} \wedge \mathrm{co}\mathcal{G}$. Then, $\mathrm{NPMV}(k)$ is the class of partial multivalued functions defined in the following way:

$$\begin{aligned} \mathrm{NPMV}(1) &= \mathrm{NPMV}, \text{ and, for } k \geq 2, \\ \mathrm{NPMV}(k) &= \mathrm{NPMV} - \mathrm{NPMV}(k-1). \end{aligned}$$

Fenner *et al.* prove that for every $k \geq 1$, $f \in \mathrm{NPMV}(k)$ if and only if f is polynomially length-bounded and $\mathrm{graph}(f) \in \mathrm{NP}(k)$.

In particular, we are interested in the class coNPMV. It follows that a function f belongs to coNPMV if and only if it is polynomially length-bounded and $\mathrm{graph}(f)$ belongs to coNP. Observe that the classes NPMV and coNPMV satisfy the nice symmetry that graphs of functions in the former class are in NP and those in the latter class are in coNP.

Just as the definition of the Boolean hierarchy over NP leads to the class $\mathrm{NP}(n^{O(1)})$ (see [Wag90]), we now introduce the class $\mathrm{NPMV}(n^{O(1)})$. It can be shown that a function h belongs to $\mathrm{NPMV}(k)$ if and only if there is a 2-ary function $f \in \mathrm{NPMV}$ such that

$$\begin{aligned}\text{set-}h(x) = \text{set-}f(x,k) - \big(&\text{set-}f(x,k-1)\\ -\big(&\text{set-}f(x,k-2)\\ -\big(\cdots -\ &\text{set-}f(x,1)\cdots\big)\big)\big).\end{aligned}$$

We say that $h \in \mathrm{NPMV}(n^{O(1)})$ if and only if there is a function $f \in \mathrm{NPMV}$ and a polynomial p such that

$$\begin{aligned}\text{set-}h(x) = \text{set-}f(x,p(|x|))\\ -\big(\text{set-}f(x,p(|x|)-1)\\ -\big(\text{set-}f(x,p(|x|)-2)\\ -\big(\cdots -\ \text{set-}f(x,1)\cdots\big)\big)\big).\end{aligned}$$

The above mentioned result by Fenner *et al.* can be extended to show that $f \in \mathrm{NPMV}(n^{O(1)})$ if and only if f is polynomially length-bounded and $\mathrm{graph}(f) \in \mathrm{NP}(n^{O(1)})$.

The primary new contribution of Fenner *et al.* is the development of hierarchies of classes of functions that access classes of partial functions as oracles. This development is based on the following description of oracle Turing machines with oracles that compute partial functions. Assume first that the oracle is a single-valued partial function g. Let $\perp$ be a symbol not belonging to the finite alphabet Σ. In order for a machine M to access a partial function oracle, M has a write-only input oracle tape, a separate read-only output oracle tape, and a special oracle call state q. To query g on a string x, M enters state q with x on the oracle input tape in the usual fashion. The oracle then returns the value $g(x)$ on the oracle output tape if the value exists, and writes $\perp$ on the tape otherwise. (It is possible that M may read only a portion of the oracle's output if the oracle's output is too long to read with the resources of M.) We shall assume, without loss of generality, that M never makes the same oracle query more than once on any possible computation path.

If g is a single-valued partial function and M is a deterministic oracle transducer as just described, then we let $M[g]$ denote the single-valued partial function computed by M with oracle g.

2.1 Definition. *[FHOS93] Let f and g be multivalued partial functions. f is* Turing reducible *to g in polynomial time, $f \leq^{\mathrm{P}}_{\mathrm{T}} g$, if for some deterministic oracle transducer M, for every single-valued refinement g' of g, $M[g']$ is a single-valued refinement of f.*

Fenner *et al.* prove that $\leq^{\mathrm{P}}_{\mathrm{T}}$ is a reflexive and transitive relation over the class of all partial multivalued functions.

Let $\mathcal{F}$ be a class of partial multivalued functions. $\mathrm{FP}^{\mathcal{F}}$ denotes the class of partial multivalued functions f that are $\leq^{\mathrm{P}}_{\mathrm{T}}$-reducible to some $g \in \mathcal{F}$. $\mathrm{FP}^{\mathcal{F}[k]}$ (respectively, $\mathrm{FP}^{\mathcal{F}[\log]}$) denotes the class of partial multivalued functions f that are $\leq^{\mathrm{P}}_{\mathrm{T}}$-reducible to some $g \in \mathcal{F}$ via a machine that, on input x, makes k adaptive queries (respectively, $\mathcal{O}(\log |x|)$ adaptive queries) to its oracle.

This definition template defines classes of multivalued partial functions such as $\mathrm{FP}^{\mathrm{NPMV}}$, and can easily be extended to define $\mathrm{NPMV}^{\mathrm{NPMV}}$. If $\mathcal{K}$ is a class of sets, then $\mathrm{FP}^{\mathcal{K}}$ is defined as usual, except that we allow it to compute partial functions (at the discretion of the oracle machine).

We will use the following generalization of the many-one metric reducibility of Krentel [Kre88] in order to discuss complete functions for classes of multivalued functions.

2.2 Definition. *Given partial multivalued functions $f, g : \Sigma^* \mapsto \Sigma^*$, we say f* is metric many-one reducible *to g, or symbolically, $f \leq^{\mathrm{P}}_{m} g$, if there are functions $t_1, t_2 \in \mathrm{FP}$ such that the multivalued partial function h defined by*

$$h(x) = t_2(x, (g \circ t_1)(x))$$

is a refinement of f, where set-$h(x)$ *is defined as*

$$\{\, t_2(x,y) \mid g(t_1(x)) \mapsto y \,\}.$$

If, in addition, we have set-$t_2(x,(g \circ t_1)(x)) =$ set-$f(x)$, *we call it a* strong metric many-one reduction, *denoted by $f \leq^{\mathrm{P}}_{sm} g$.*

The motivation underlying this definition is that, given a value of $g(x)$, one can compute in polynomial time a value of $f(x)$. In the case of a strong reduction, one gets all values of $f(x)$ when varying over all values of $g(x)$. Obviously, $f \leq^{\mathrm{P}}_{m} g$ implies $f \leq^{\mathrm{P}}_{\mathrm{T}} g$.

The classes that we have been considering relate in interesting ways to studies of the complexity of optimization problems. In order to capture the complexity of optimization problems, Krentel [Kre88] defined the complexity classes MaxP and MinP as the functions computable by taking the maximum, respectively minimum, over sets of feasible solutions of problems in NP. Further, Krentel extended these classes to hierarchies of classes of optimization functions [Kre92]. Krentel defined these functions using the notion of a *metric Turing machine*, which we now review. Consider nondeterministic polynomial time Turing machines that print an output value on every path. We associate with every inner node of the computation tree either the function min or the function max (for the classes MinP and MaxP, resp.; all nodes are associated with the same function), Thus, metric Turing machines define (total) functions from input words to integers via the usual bottom-up evaluation of the machine's computation tree. Since all the function classes considered in this paper are partial, we extend the metric Turing machine just defined by allowing the machine to output a special symbol $\perp$ which denotes that the computation on the corresponding path ends with an undefined result. We extend the min and max functions in

the obvious way: define $\max(x,\bot) = \max(\bot,x) = x$ and $\min(x,\bot) = \min(\bot,x) = x$, for all x (including $\bot$ itself). Vollmer and Wagner [VW93, VW95] gave a detailed structural examination of Krentel's hierarchy. Here, we just define the class MaxP using an operator-characterization from [VW95]. MinP is defined analogously.

$$h \in \text{MaxP} \iff \exists f,\ g \in \text{FP}:\ h(x) = \max_{0 \le y \le g(x)} f(x,y).$$

3. Functions Complete for coNPMV

NPMV is precisely the class of functions that compute witnesses for NP sets in the following sense: For any set $L \in \text{NP}$ there exist a set $A \in \text{P}$ and a polynomial p such that for all x, we have

$$x \in L \iff \exists y \in \Sigma^{p(|x|)}\colon (x,y) \in A.$$

Any y such that $(x,y) \in A$ is called a *witness for* x (with respect to A). Clearly, there is a function $f \in \text{NPMV}$ such that set-$f(x)$ is exactly the set of witnesses for x. On the other hand, any NPMV function f defines a set in NP as the set of all x such that there exists an output of f on x. In other words, $\text{dom}(f) \in \text{NP}$. As a consequence of this discussion, we see (abusing notation) that $\text{dom}(\text{NPMV}) = \text{NP}$.

Next, we extend the notion of a witness to Σ_2^p. For any Σ_2^p set L there exist a set $B \in \text{coNP}$ and a polynomial p such that for all x, we have $x \in L \iff \exists y \in \Sigma^{p(|x|)}\colon (x,y) \in B$. A y such that $(x,y) \in B$ is called a *witness for* x (with respect to B). What function class captures the computation of witnesses for Σ_2^p sets? Since $\Sigma_2^p = \text{NP}^{\text{NP}}$, certainly witnesses can be computed in NPMV^{NP}. However, we will see below that the seemingly weaker class coNPMV already suffices to do so.

Let us consider set L again. We may safely assume that for all $(x,y) \in B$ we have $y \in \Sigma^{p(|x|)}$. Since $B \in \text{coNP}$, it is then the graph of a coNPMV function f, so that set-$f(x)$ is exactly the set of witnesses for x. Hence, coNPMV can compute witnesses for sets in Σ_2^p. Conversely, for any coNPMV function f, we have $\text{dom}(f) \in \Sigma_2^p$. This is because, for any x, $x \in \text{dom}(f) \iff \exists y \in \Sigma^{p(|x|)}\colon y \in \text{set-}f(x)$. Thus, coNPMV is *precisely* the class of functions that computes witnesses for Σ_2^p sets. As a consequence, we have the following proposition.

3.1 Proposition. $\text{dom}(\text{coNPMV}) = \Sigma_2^p$.

Witnesses of Σ_2^p complete sets can give rise to complete functions for coNPMV. Consider, for example, the satisfiability problem QBF_2 for Boolean formulas with two quantifiers. Let φ be a Boolean formula in the variables $\mathbf{x} = (x_1,\ldots,x_k)$ and $\mathbf{y} = (y_1,\ldots,y_l)$. Then we define

$$\varphi(\mathbf{x},\mathbf{y}) \in \text{QBF}_2 \iff \exists \mathbf{x}\ \forall \mathbf{y}\colon \varphi(\mathbf{x},\mathbf{y}) = 1.$$

Let F_2 be the multivalued function that computes witnesses, i.e., partial assignments $\mathbf{x} = (x_1,\ldots,x_k)$, for QBF_2 formulas φ as above.

3.2 Theorem. *F_2 is $\le_m^{\text{P}}$-complete for* coNPMV.

Proof. We have argued already that $F_2 \in \text{coNPMV}$. Let f be any coNPMV function. There is an NP transducer M and a polynomial p such that for all x, we have set-$f(x) = \Sigma^{p(|x|)} - \text{set-}M(x)$. We show how to compute a $y \in \text{set-}f(x)$ from $F_2(\varphi_x)$, for an appropriately constructed formula φ_x.

Define a machine M' on input x as follows. First, M' guesses a $y \in \Sigma^{p(|x|)}$. Then, M' simulates M on input x. If M outputs y on the simulated path, then M' rejects. Otherwise, M' accepts.

We have to define the reduction functions t_1 and t_2 as required in Definition 2.2. Function t_1 is Cook's reduction applied to x with M' as the underlying machine. This will give a Boolean formula φ_x that, intuitively, describes the work of M' on input x. The variables of φ_x can be partitioned into two parts:

- say $y_1,\ldots,y_k$, that are used to describe that M' guesses a $y \in \Sigma^{p(|x|)}$, and
- say $z_1,\ldots,z_l$, that are used to describe the subsequent simulation of M.

Furthermore, from any setting of the variables $y_1,\ldots,y_k$ of φ_x, we can reconstruct in polynomial time the $y \in \Sigma^{p(|x|)}$ guessed by M'. This is done by function t_2.

Let us fix a setting of the variables $y_1,\ldots,y_k$ and let $y \in \Sigma^{p(|x|)}$ be the corresponding string guessed by M'. Then we have

$$\begin{aligned} &\forall z_1,\ldots,z_l:\ \varphi_x(y_1,\ldots,y_k,z_1,\ldots,z_l) = 1 \\ &\iff\ M' \text{ accepts on all paths following } y \\ &\iff\ y \notin \text{set-}M(x) \\ &\iff\ f(x) \mapsto y, \end{aligned}$$

and hence, $t_2(x, F_2 \circ t_1(x))$ computes a refinement of $f(x)$. □

A crucial point in the above proof is that Cook's reduction maintains witnesses. That is, from a given assignment for the constructed formula φ_x one can recover a corresponding path of the nondeterministic machine. Thus any Σ_2^p complete set sharing this property with QBF_2 defines a coNPMV complete function in an analogous way. Note moreover that there is a one-one correspondence between witnesses and assignments (that is computable and invertible in polynomial time), so that the above proof in fact shows that F_2 is complete for coNPMV under $\le_{sm}^{\text{P}}$-reductions.

As an example, consider the following set L_f. For any NPMV function f and some polynomial p such that f maps strings of length n to strings of length $p(n)$, $x \in L_f$ if and only if

$$\exists y \in \Sigma^{p(|x|)/2}\ \forall z \in \Sigma^{p(|x|)/2}:\ f(x) \not\mapsto yz.$$

In other words, string y is not a prefix of an output of $f(x)$.

Clearly, for every $f \in$ NPMV, we have that L_f is in Σ_2. Thus, in particular, taking $f = sat$, L_{sat} is in Σ_2^p, is even Σ_2^p complete and has the above mentioned property. We conclude that the corresponding witness function, $not - pre - sat$, is complete for coNPMV, where

$$not - pre - sat(\varphi) \mapsto y \iff$$
y is a truth assignment of the first half of φ's variables which is not a prefix of a satisfying assignment of φ.

3.3 Theorem. *$not - pre - sat$ is $\leq_m^{\mathrm{P}}$-complete for the class* coNPMV.

Theorem 3.3 can be seen directly via a straightforward metric reduction from F_2.

4. Properties of coNPMV

NPMV is closed under $\leq_{sm}^{\mathrm{P}}$-reductions, but not under $\leq_m^{\mathrm{P}}$-reductions; in fact, it is possible to have $g \in$ NPMV and $f \leq_m^{\mathrm{P}} g$ but graph(f) be noncomputable. (For example, define f to map x to two values, one of them solves the halting problem on x, i.e., is from $\{0,1\}$, the second value is constant 10. Then clearly f is not recursive, but the constant function 10 is a refinement of f in NPMV.) However, NPMV is closed under this reduction in a weaker sense, defined below.

4.1 Definition. A class $\mathcal{F}$ is *c-closed* under reducibility $\leq_r$, if $g \in \mathcal{F}$ and $f \leq_r g$ implies $f \in_c \mathcal{F}$.

It is immediate from this definition that NPMV is c-closed under $\leq_m^{\mathrm{P}}$-reductions. One might suspect that this same fact holds for coNPMV. However, it is quite unlikely that coNPMV is c-closed under this reducibility: otherwise, since $sat \in$ coNPMV and sat is complete for NPMV, we would get that NPMV $\subseteq_c$ coNPMV. But this seems to be very unlikely as the following extension of a result of Fenner *et al.* [FHOS93] shows.

4.2 Theorem. NPMV $\subseteq$ coNPMV $\iff$ NPMV $\subseteq_c$ coNPMV $\iff$ NP = coNP.

Proof. We cycle through the implications. The first implication is trivial. For the second, let $L \in$ NP. Define function

$$\chi_L(x) = \begin{cases} 1 & \text{if } x \in L \\ \perp & \text{otherwise .} \end{cases}$$

Then we have $\chi_L \in$ NPMV, and hence, by assumption, $\chi_L \in$ coNPMV. Therefore, graph$(\chi_L) \in$ coNP, which implies that $L \in$ coNP since $x \in L \iff (x,1) \in$ graph(χ_L).

Now suppose that NP = coNP and let $f \in$ NPMV. Then graph$(f) \in$ NP, and therefore in coNP by assumption. Thus $f \in$ coNPMV.

□

4.3 Corollary. coNPMV *is c-closed under $\leq_m^{\mathrm{P}}$-reducibility if and only if* NP = coNP.

We observe that the proof of Theorem 4.2 shows also that NPSV $\subseteq$ coNPMV $\iff$ NP = coNP, even though it is fairly easy to see that NPSV$_t$, the class of all total NPSV functions, *is* contained in coNPMV. We also note that Theorem 4.2 extends to higher levels of the difference hierarchies over NPMV and NP, that is, NPMV$(k) \subseteq$ coNPMV$(k) \iff$ NPMV$(k) \subseteq_c$ coNPMV$(k) \iff$ NP(k) = coNP(k). By a result of Kadin [Kad88], a collapse of the Boolean hierarchy implies a collapse of the polynomial time hierarchy. Hence, there is a whole hierarchy between coNPMV and NPMV$^{\mathrm{NP}}$.

4.4 Theorem. *For all $k \geq 2$, we have*

$$\text{coNPMV} \subseteq \text{NPMV}(2) \subseteq \text{NPMV}(k) \subseteq$$
$$\text{NPMV}(k+1) \subseteq \text{NPMV}(n^{O(1)}) \subseteq \text{NPMV}^{\mathrm{NP}}.$$

Furthermore, all of the inclusions are strict unless the polynomial time hierarchy collapses.

Proof. It remains to show the last inclusion. Let $f \in$ NPMV$(n^{O(1)})$. Then the graph of f is in NP$(n^{O(1)})$, which is known to be equal to P$^{\mathrm{NP[log]}}$ [Wag90]. Obviously, f can be computed by an NPMV algorithm with access to a P$^{\mathrm{NP[log]}}$ oracle: simply guess an output of f and querying its graph check that the guess is correct. Thus, NPMV$(n^{O(1)}) \subseteq$ NPMV$^{\mathrm{P^{NP[log]}}} \subseteq$ NPMV$^{\mathrm{NP}}$.

□

It follows from Theorem 4.2 that we can have the situation that a function that is complete for some class, like *sat* for NPMV, is in coNPMV without the corresponding class being contained in coNPMV. This seems to happen again for *maxsat*, the function that maps a Boolean formula to its lexicographically largest satisfying assignment. Fenner *et al.* [FHOS93] show that *maxsat* $\in$ NPMV(2). In fact, it is even in coNPMV. However, we will show that the corresponding classes, namely MaxP or FP$^{\mathrm{NP}}$ are unlikely to be contained in coNPMV.

4.5 Theorem. *maxsat* $\in$ coNPMV.

Proof. Consider an NPMV machine M that, on input of a formula φ, guesses an assignment y for φ. If y does not satisfy φ, then M outputs y. Otherwise, if y does satisfy φ, M guesses another assignment $y' > y$. If y' also satisfies φ, M outputs y, otherwise M rejects.

M outputs every assignment except the maximum satisfying one (if there is one). Hence *maxsat* $\in$ coNPMV. □

Krentel [Kre92] showed that FP$^{\mathrm{NP}}$ = FP$^{\mathrm{MaxP[1]}}$. Since FP$^{\mathrm{NPMV}}$ = FP$^{\mathrm{NP}}$ [FHOS93] and *maxsat* is complete for MaxP, we have that FP$^{\mathrm{NPMV}} \subseteq$ FP$^{\mathrm{coNPMV[1]}}$. That is, polynomially many queries of a

FP function to NPMV can be replaced by one query to coNPMV. Hence, as we have mentioned, coNPMV seems to be a more powerful class than NPMV. We will give more evidence for this in the next section.

It is tempting to conclude from the proof of Theorem 4.5 that $\mathrm{Max\,P} \subseteq \mathrm{coNPMV}$. However, this is unlikely.

4.6 Corollary. $\mathrm{Max\,P} \subseteq \mathrm{coNPMV} \iff \mathrm{Min\,P} \subseteq \mathrm{coNPMV} \iff \mathrm{NP} = \mathrm{coNP}$.

Proof. If $\mathrm{Max\,P} \subseteq \mathrm{coNPMV}$, then $\mathrm{NPMV} \subseteq_c \mathrm{Max\,P} \subseteq \mathrm{coNPMV}$, and therefore $\mathrm{NPMV} \subseteq_c \mathrm{coNPMV}$. But by Theorem 4.2, this implies $\mathrm{NP} = \mathrm{coNP}$. Conversely, if $\mathrm{NP} = \mathrm{coNP}$, then $\mathrm{NPMV}^{\mathrm{NP}} = \mathrm{NPMV}^{\mathrm{NP} \cap \mathrm{coNP}} = \mathrm{NPMV}$. This implies $\mathrm{MaxP} \subseteq \mathrm{NPMV}$, and since the hypothesis also implies $\mathrm{NPMV} = \mathrm{coNPMV}$, that $\mathrm{MaxP} \subseteq \mathrm{coNPMV}$. □

We conclude this section with an observation regarding the relationship between MaxP and NPMV. First, note that trivially $\mathrm{NPSV} \subseteq \mathrm{MaxP} \cap \mathrm{MinP}$, since *the* output of an NPSV function is both the minimum and the maximum. Similarly, $\mathrm{NPMV} \subseteq_c \mathrm{MaxP} \cap \mathrm{MinP}$. The more interesting question is whether these inclusions are strict. This is quite likely.

4.7 Theorem. $\mathrm{Max\,P} \subseteq \mathrm{NPMV} \iff \mathrm{Min\,P} \subseteq \mathrm{NPMV} \iff \mathrm{NP} = \mathrm{coNP}$.

Proof. If $\mathrm{NP} = \mathrm{coNP}$, then $\mathrm{FP}^{\mathrm{NP}} \subseteq \mathrm{NPMV}$ [Sel94], thus especially $\mathrm{Max\,P} \cup \mathrm{Min\,P} \subseteq \mathrm{NPMV}$.

Now suppose $\mathrm{Max\,P} \subseteq \mathrm{NPMV}$ (the case for Min P is analogous). Let $L \in \mathrm{coNP}$. Define

$$f(x) = \begin{cases} 0 & \text{if } x \in L \\ 1 & \text{otherwise.} \end{cases}$$

Then $f \in \mathrm{Max\,P}$ and hence, by assumption, in NPMV. Since $x \in L$ if and only if $f(x) = 0$, we have $L \in \mathrm{NP}$. □

The last two results relativize: analogous results hold for higher levels of the NPMV hierarchy and Krentel's min/max hierarchy [FHOS93, VW95]. For the relativized version of Theorem 4.7 one has to use techniques from Krentel [Kre92] and Vollmer and Wagner [VW95].

5. A Characterization of coNPMV

As we have already seen in the preceding section, coNPMV seems to be a more powerful class than NPMV. This is somewhat surprising in light of the aforementioned symmetry in the definitions of coNPMV and NPMV by their graphs.

The following theorem shows that coNPMV is in fact very close to $\mathrm{NPMV}^{\mathrm{NP}}$. This is surprising as well, as we have already seen in Corollary 4.4 that there is a hierachy of function classes between coNPMV and $\mathrm{NPMV}^{\mathrm{NP}}$.

If f is a multivalued function and g is a single valued function, then $g \circ f$ is defined by $\mathrm{graph}(g \circ f) = \{ \langle x, g(y) \rangle \mid f(x) \mapsto y \}$. Let π_2^1 denote the projection function that maps a pair of strings to its first component. By $\pi_2^1 \circ \mathrm{coNPMV}$ we denote $\{ \pi_2^1 \circ f \mid f \in \mathrm{coNPMV} \}$.

5.1 Theorem. $\mathrm{NPMV}^{\mathrm{NP}} = \pi_2^1 \circ \mathrm{coNPMV}$.

Proof. The right to left containment follows from Corollary 4.4 and the fact that the projection of any $\mathrm{NPMV}^{\mathrm{NP}}$ function is still in $\mathrm{NPMV}^{\mathrm{NP}}$, hence $\pi_2^1 \circ \mathrm{coNPMV} \subseteq \mathrm{NPMV}^{\mathrm{NP}}$.

For the other direction, let $f \in \mathrm{NPMV}^{\mathrm{NP}}$. By a standard argument there is a polynomial q and a predicate $R \in \mathrm{P}$ such that for any x and $y \in \Sigma^{q(|x|)}$

$$f(x) \mapsto y \iff \exists z \in \Sigma^{q(|x|)}\ \forall w \in \Sigma^{q(|x|)} \colon R(x, y, z, w).$$

(Informally, z witnesses that there is a path outputting y where all queries answered affirmatively are legitimate, and w is used to verify the negatively answered queries.) Define f' such that for any x and y, $z \in \Sigma^{q(|x|)}$

$$f'(x) \mapsto \langle y, z \rangle \iff \forall w \in \Sigma^{q(|x|)} \colon R(x, y, z, w).$$

So $\neg R$ witnesses that $f' \in \mathrm{coNPMV}$. But $f(x) = \pi_2^1 \circ f'(x)$, which shows that $f \in \pi_2^1 \circ \mathrm{coNPMV}$. □

There is a counterintuitive facet to Theorem 5.1. The reason why it is likely that coNPMV is a proper subclass of $\mathrm{NPMV}^{\mathrm{NP}}$ is not because outputs of coNPMV functions give too little information, but rather that they give too much. We can compute an arbitrary $\mathrm{NPMV}^{\mathrm{NP}}$ function simply by throwing away part of the output of a coNPMV function. This is what the projection operator accomplishes, and it is most likely necessary.

Applying Theorem 5.1, many properties of $\mathrm{NPMV}^{\mathrm{NP}}$ now carry over to coNPMV. In Section 3 we have shown functions F_2 and $not - pre - sat$ complete for coNPMV. Since the projection function is in FP, we get that those functions are complete for $\mathrm{NPMV}^{\mathrm{NP}}$ as well.

5.2 Corollary. $\mathrm{NPMV}^{\mathrm{NP}}$ *is the c-closure of* coNPMV *under* $\leq_m^{\mathrm{P}}$*-reducibility and the closure of* coNPMV *under* $\leq_{sm}^{\mathrm{P}}$*-reducibility.*

In particular, we get

5.3 Corollary. $\mathrm{FP}^{\mathrm{coNPMV[1]}} = \mathrm{FP}^{\mathrm{NPMV}^{\mathrm{NP}}[1]}$ *and* $\mathrm{FP}^{\mathrm{coNPMV}} = \mathrm{FP}^{\mathrm{NPMV}^{\mathrm{NP}}} = \mathrm{FP}^{\mathrm{NP}^{\mathrm{NP}}}$.

Observe by contrast that $\mathrm{FP}^{\mathrm{NPMV}} = \mathrm{FP}^{\mathrm{NP}} = \mathrm{FP}^{\mathrm{Min\,P}} = \mathrm{FP}^{\mathrm{Max\,P}}$, so coNPMV and NPMV define different Δ-levels of the functional polynomial hierarchy.

Fenner *et al.* [FHOS93] have shown that $\mathrm{NPMV}(2) \subseteq \mathrm{FP}^{\mathrm{NPMV}} \Longleftrightarrow \Sigma_2^p = \Delta_2^p$. Note that in contrast for the corresponding language classes we have $\mathrm{NP}(k) \subseteq \mathrm{P}^{\mathrm{NP}}$ for all k. We can now improve the result of Fenner *et al.*

5.4 Corollary. $\mathrm{coNPMV} \subseteq \mathrm{FP}^{\mathrm{NPMV}} \Longleftrightarrow \Sigma_2^p = \Delta_2^p$.

Proof. If $\Sigma_2^p = \Delta_2^p$, then

$$\mathrm{coNPMV} \subseteq \mathrm{FP}^{\mathrm{coNPMV}} = \mathrm{FP}^{\mathrm{NPMV}^{\mathrm{NP}}} = \mathrm{FP}^{\Sigma_2^p} = \mathrm{FP}^{\Delta_2^p} = \mathrm{FP}^{\mathrm{NP}} = \mathrm{FP}^{\mathrm{NPMV}},$$

where the last equality is Theorem 1 in [FHOS93] and the second follows from the relativized version of the same theorem. Conversely, if $\mathrm{coNPMV} \subseteq \mathrm{FP}^{\mathrm{NPMV}}$, then $\mathrm{dom}(\mathrm{coNPMV}) \subseteq \mathrm{P}^{\mathrm{NP}} = \Delta_2^p$, so that $\Sigma_2^p \subseteq \Delta_2^p$. □

5.5 Corollary. *For any $k \geq 1$, we have*

$$\mathrm{NPMV}^{\mathrm{NP}} \subseteq \mathrm{FP}^{\mathrm{coNPMV}[1]} \subseteq \mathrm{FP}^{\mathrm{coNPMV}[k]} \subseteq$$

$$\mathrm{FP}^{\mathrm{coNPMV}[k+1]} \subseteq \mathrm{FP}^{\mathrm{coNPMV}} = \mathrm{FP}^{\mathrm{NP}^{\mathrm{NP}}}.$$

Furthermore, all inclusions are strict unless the polynomial time hierarchy collapses.

Proof. It remains to show the strictness of the inclusions. Suppose $\mathrm{FP}^{\mathrm{coNPMV}[1]} \subseteq \mathrm{NPMV}^{\mathrm{NP}}$. This is equivalent to $\mathrm{FP}^{\mathrm{NPMV}^{\mathrm{NP}}[1]} \subseteq \mathrm{NPMV}^{\mathrm{NP}}$, which implies $\mathrm{P}^{\Sigma_2^p[1]} \subseteq \Sigma_2^p$. But then $\Pi_2^p = \Sigma_2^p = \mathrm{PH}$. For the other inclusions, suppose $\mathrm{FP}^{\mathrm{coNPMV}[k]} = \mathrm{FP}^{\mathrm{coNPMV}[k+1]}$. Then $\mathrm{FP}^{\mathrm{NPMV}^{\mathrm{NP}}[k]} = \mathrm{FP}^{\mathrm{NPMV}^{\mathrm{NP}}[k+1]}$. By a theorem of Fenner *et al.* [FHOS93], this implies that $\mathrm{FP}^{\Sigma_2^p[k]} = \mathrm{FP}^{\Sigma_2^p[k+1]}$, which, by a relativization of Kadin's theorem [Kad88], implies that the polynomial hierarchy collapses. □

Thus we see, combining Theorems 4.4 and 5.5, that all classes of the difference hierarchy over NPMV are included in the query hierarchy over coNPMV, in fact already in its first level. There are (under reasonable assumptions) no inclusions in the opposite direction. Concerning the relationship between the query hierarchy over NPMV and the difference hierarchy over NPMV, we know from Fenner *et al.* [FHOS93] that all classes of the first hierarchy are included in certain classes of the second hierarchy. Any inclusion in the opposite direction implies $\mathrm{coNPMV} \subseteq \mathrm{FP}^{\mathrm{NPMV}}$, which again implies a collapse of the polynomial time hierarchy, by Corollary 5.4.

6. Relationships Between the Functional Difference and Polynomial Time Hierarchies

Chang and Kadin [CK90] showed that if the Boolean hierarchy over NP collapses to the k^{th} level, then the polynomial hierarchy collapses to the k^{th} level of the Boolean hierarchy over $\mathrm{NP}^{\mathrm{NP}}$: if $\mathrm{NP}(k+1) = \mathrm{NP}(k)$, then $\mathrm{PH} = \mathrm{NP}^{\mathrm{NP}}(k)$. It is a simple consequence of known results that a similar connection exists for the corresponding functional hierarchies, namely $\mathrm{NPMV}(k)$ and $\Sigma\mathrm{MV}_k = \mathrm{NPMV}^{\Sigma_{k-1}^p}$.

6.1 Theorem. *For any $k \geq 1$, if $\mathrm{NPMV}(k+1) = \mathrm{NPMV}(k)$ then $\Sigma\mathrm{MV}_3 = \mathrm{NPMV}^{\mathrm{NP}}(k)$.*

Proof. $\mathrm{NPMV}(k+1) = \mathrm{NPMV}(k)$ is equivalent with $\mathrm{NP}(k+1) = \mathrm{NP}(k)$ [FHOS93], which implies $\Sigma_3^p = \mathrm{NP}^{\mathrm{NP}}(k)$ [CK90] (relativized). Since $\Sigma_{k+1}^p = \Sigma_k^p \Longleftrightarrow \Sigma\mathrm{MV}_{k+1} = \Sigma\mathrm{MV}_k$ [FHOS93], we get that $\Sigma\mathrm{MV}_3 = \mathrm{NPMV}^{\mathrm{NP}}(k)$. □

Since $\mathrm{NP}^{\mathrm{NP}}(k) \subseteq \mathrm{P}^{\mathrm{NP}^{\mathrm{NP}}[k]}$, a consequence of Chang and Kadin's theorem is that if $\mathrm{NP}(k+1) = \mathrm{NP}(k)$, then $\Sigma_3^p = \mathrm{P}^{\mathrm{NP}^{\mathrm{NP}}[k]}$ (indeed, they prove this directly in their paper before treating the stronger result). The functional analogue of such a collapse would be $\Sigma\mathrm{MV}_3 = \mathrm{FP}^{\mathrm{NPMV}^{\mathrm{NP}}[k]}$ or, equivalently, $\Sigma\mathrm{MV}_3 = \mathrm{FP}^{\mathrm{coNPMV}[k]}$. We cannot expect this as a direct consequence of Theorem 6.1, since the difference and query hierarchies are not intertwined in this context. Nevertheless, such an analogous result does hold. To see this, we have to modify the proof of the Chang and Kadin theorem.

6.2 Theorem. *If $\mathrm{NPMV}(k+1) = \mathrm{NPMV}(k)$ then $\Sigma\mathrm{MV}_3 = \mathrm{NPMV} \circ \mathrm{FP}^{\mathrm{coNPMV}[k-1]}$.*

Proof. (Sketch.) In order to explain how Chang and Kadin's proof gives this result, we recall some of their definitions, with some minor modifications in notation (for greater detail, we refer the reader to their paper [CK90]). Denote the $\leq_m^p$-complete language for $\mathrm{NP}(k)$ (respectively $\mathrm{coNP}(k)$) as $L_{\mathrm{NP}(k)}$ (respectively $L_{\mathrm{coNP}(k)}$). For example, $L_{\mathrm{NP}(1)} = \mathrm{SAT}$ and $L_{\mathrm{NP}(2)} = \{\langle x_1, x_2\rangle \mid x_1 \in \mathrm{SAT} \text{ and } x_2 \in \overline{\mathrm{SAT}}\}$. Since, by hypothesis, $\mathrm{NP}(k) = \mathrm{coNP}(k)$, it follows that $L_{\mathrm{NP}(k)} \leq_m^p L_{\mathrm{coNP}(k)}$. The basic idea underlying the Chang and Kadin proof is that such a reduction induces a reduction from an initial segment of SAT to an initial segment of $\overline{\mathrm{SAT}}$. This is done inductively via the notion of a "hard sequence", which is a j-tuple which, together with a $\leq_m^p$-reduction from $\mathrm{NP}(k)$ to $\mathrm{coNP}(k)$, can be used to find a $\leq_m^p$-reduction from $\mathrm{NP}(k-j)$ to $\mathrm{coNP}(k-j)$.

6.3 Definition. Let $L_{\mathrm{NP}(k)} \leq_m^p L_{\mathrm{coNP}(k)}$ via some polynomial time function h. Then we call

$\langle 1^m, x_1, \ldots, x_j \rangle$ a *hard sequence* with respect to h for length m of order j, if either $j = 0$ or the following conditions hold:

1. $1 \leq j \leq k-1$,
2. $|x_j| \leq m$,
3. $x_j \in \overline{\mathrm{SAT}}$,
4. $\langle 1^m, x_1, \ldots, x_{j-1} \rangle$ is a hard sequence with respect to h, and
5. for all $y_1, \ldots, y_\ell \in \Sigma^{\leq m}$ (where $\ell = k - j$), $\pi_{\ell+1} \circ h(\langle y_1, \ldots, y_\ell, x_j, \ldots, x_1 \rangle) \in \overline{\mathrm{SAT}}$.

A hard sequence is called *maximal* if it cannot be extended to a hard sequence of a higher order. In this case the order of the sequence j is said to be maximal.

We can now outline the proof. Chang and Kadin's Lemma 3 [CK90] states that, given a maximal hard sequence for an appropriate (polynomially bounded) length, an NP machine can recognize an initial segment of the canonical complete language for $\mathrm{NP}^{\mathrm{NP}}$. That is, with the aid of such a sequence we can replace a Σ_2^p machine with an NP machine. Thus it suffices to find a maximal hard sequence to collapse the NP's of a $\Sigma\mathrm{MV}_3 = \mathrm{NPMV}^{\Sigma_2^p}$ machine.

Our principle observation is this: *Hard sequences of any given order can be obtained by a single query to a* coNPMV *oracle.* This can easily be seen as follows. Define the function $H\colon 1^+ \times \mathbf{N} \mapsto \Sigma^*$ such that $H(1^m, j) \mapsto \langle 1^m, x_1, \ldots, x_j \rangle$ if and only if $\langle 1^m, x_1, \ldots, x_j \rangle$ is a hard sequence for length m of order j. It follows from Definition 6.3 that the set of hard sequences is in coNP [CK90]; hence, $\mathrm{graph}(H) \in$ coNP, so that $H \in$ coNPMV. Therefore, we can obtain a maximal hard sequence for the appropriate polynomial length $m = p(|x|)$ by querying a coNPMV oracle for the value of $H(m, j)$ for j varying from 1 to $k-1$. We then feed the resulting maximal hard sequence, along with the original input x, to an NPMV machine which can, via the induced reduction from coNP to NP, collapse the NP oracles in an $\mathrm{NPMV}^{\mathrm{NP}^{\mathrm{NP}}}$ computation. □

7. A Remark on Counting Classes

The results of our paper show that in the context of relational structures computed by polynomial time machines, in a sense the universal mode is more powerful than the existential one. In the context of counting classes, a similar phenomenon has been observed by Seinosuke Toda [Tod91]. In this section, we briefly show that Toda's result is a special case of one of our observations.

Recall the following general definition of counting classes from [Tod91, VW93]:

Let $\mathcal{K}$ be a class of sets. Then, $\# \cdot \mathcal{K}$ consists of those functions f for which there exist a set $A \in \mathcal{K}$ and a polynomial p such that for all x,

$$f(x) = \left| \{ y \mid |y| \leq p(|x|) \wedge \langle x, y \rangle \in A \} \right|.$$

It is obvious that $\# \cdot \mathrm{P} = \#\mathrm{P}$, and moreover it can be shown that $\# \cdot \mathrm{NP} = \mathrm{spanP}$.

We have the following relationship to our classes of functions:

7.1 Proposition.

1. $\# \cdot \mathrm{NP}$ *consists of exactly those functions h for which there exists a function $f \in$ NPMV such that for all x, $h(x) = \left|\text{set-}f(x)\right|$.*
2. $\# \cdot \mathrm{coNP}$ *consists of exactly those functions h for which there exists a function $f \in$ coNPMV such that for all x, $h(x) = \left|\text{set-}f(x)\right|$.*

Now we have the following surprising result, which was already proved by Toda [Tod91, Theorem 4.1.6]:

7.2 Corollary. $\# \cdot \mathrm{coNP} = \#\mathrm{P}^{\mathrm{NP}}$.

Proof. Immediate by the preceding proposition and the fact that coNPMV is the class of functions that compute witnesses for Σ_2^p computations, see Proposition 3.1. □

Acknowledgment

We appreciate the contributions of J. Ramachandran to the results presented here.

References

[BLS84] R. Book and T. Long and A. Selman Quantitative relativizations of complexity classes *SIAM J. on Comput.*, 13(3):461–487, August 1984.

[CK90] R. Chang and J. Kadin. The boolean hierarchy and the polynomial hierarchy: a closer connection. In *Proceedings of the 5th Annual Conf. on Structure in Complexity Theory*, pages 140–153, New York, 1990. IEEE Computer Society Press.

[FHOS93] S. Fenner, S. Homer, M. Ogihara, and A. Selman. On using oracles that compute values. In *Proc. 10th Annual Symp. on Theoret. Aspects of Computer Science, Lecture Notes in Computer Science*, volume 665, pages 398–407. Springer-Verlag, 1993.

[Kad88] J. Kadin. The polynomial time hierarchy collapses if the Boolean hierarchy collapses. *SIAM J. on Comput.*, 17(6):1263–1282, December 1988.

[Kre88] M. Krentel. The complexity of optimization problems. *J. Comput. System Sci.*, 36:490–509, 1988.

[Kre92] M. Krentel. Generalizations of OptP to the polynomial hierarchy. *Theor. Com. Sci.*, 97:183–198, 1992.

[Pap84] C. Papadimitriou. On the complexity of unique solutions. *J. of the ACM*, 31:392–400, 1984.

[Ram94] J. Ramachandran. The polynomial time function hierarchy. Technical Report no. 94-50, Department of Computer and Information Sciences, Ohio State Univeristy, 1994.

[Sel94] A. Selman. A taxonomy of complexity classes of functions. *J. Comput. System Sci.*, 48(2):357–381, 1994.

[Tod91] S. Toda. Computational Complexity of Counting Complexity Classes. PhD thesis, Tokyo Institute of Technology, Department of Computer Science, Tokyo, Japan, 1991.

[VW93] H. Vollmer and K. Wagner. The complexity of finding middle elements. *International Journal of Foundations of Computer Science*, 4:293–307, 1993.

[VW95] H. Vollmer and K. Wagner. Complexity classes of optimization functions. *Information and Computation*, 120:198–219, 1995.

[Wag87] K. Wagner. More complicated questions about maxima and minima, and some closures of NP. *Theoret. Comput. Sci.*, 51:53–80, 1987.

[Wag90] K. Wagner. Bounded query classes. *SIAM J. Comput.*, 19:833–846, 1990.