

A polynomial-time approximation algorithm for a geometric dispersion problem

Marc Benkert^{1*}, Joachim Gudmundsson², Christian Knauer³, Esther Moet⁴,
René van Oostrum⁴, and Alexander Wolff^{1*}

¹ Faculty of Computer Science, Karlsruhe University, P.O. Box 6980,
D-76128 Karlsruhe, Germany. <http://i11www.iti.uka.de/algo/group>

² National ICT Australia Ltd[‡], Sydney, Australia,
joachim.gudmundsson@nicta.com.au

³ Institute of Computer Science, Freie Universität Berlin, Germany.
christian.knauer@inf.fu-berlin.de

⁴ Department of Computing and Information Sciences, Universiteit Utrecht,
The Netherlands. {esther,rene}@cs.uu.nl

Abstract. We consider the problem of placing a maximal number of disks in a rectangular region containing obstacles such that no two disks intersect. Let α be a fixed real in $(0, 1]$. We are given a bounding rectangle P and a set \mathcal{R} of possibly intersecting unit disks whose centers lie in P . The task is to pack a set \mathcal{B} of m disjoint disks of radius α into P such that no disk in \mathcal{B} intersects a disk in \mathcal{R} , where m is the maximum number of *unit* disks that can be packed. Baur and Fekete showed that the problem cannot be solved in polynomial time for $\alpha \geq 13/14$, unless $\mathcal{P} = \mathcal{NP}$. In this paper we present an algorithm for $\alpha = 2/3$.

1 Introduction

Obnoxious facility location problems consider the placement of facilities of which clients consider it undesirable to be in the proximity, for instance, nuclear power plants or garbage dumps. There are several models for and variations to the problem; see the survey by Cappanera [3]. We consider the following instance: we are given a bounding rectangle P , a set \mathcal{R} of n points in P (the red points), and an integer k , and we should construct a set \mathcal{B} of k (blue) points such that the minimum distance from a blue point to another point (either red or blue) is maximized over all points in \mathcal{B} . If the optimal distance is denoted by r_{opt} , then we can reformulate the problem as follows: we are given a set of n centers of possibly overlapping red disks with unknown radius r_{opt} , and we are to determine r_{opt} and to find a set of k blue disks with radius r_{opt} such that no blue disk overlaps any other disk, whether red or blue.

* Supported by grant WO 758/4-1 of the German Science Foundation (DFG). Part of this work was done while M. Benkert visited NICTA on a DAAD grant.

‡ Funded by the Australian Government's Backing Australia's Ability initiative, in part through the Australian Research Council.

The problem of packing objects into a bounded region is one of the classic problems in mathematics and theoretical computer science, see for example the monographs [11, 13] which are solely devoted to this problem, and the survey by Tóth [12]. In this paper we consider problems related to packing disks into a polygonal region. As pointed out by Baur and Fekete in [2], even when the structure of the region and the objects are simple, only very little is known, see for example [6, 9].

Consider the following decision problem corresponding to our optimization problem: we are given \mathcal{R} and k and a radius r , and we must decide whether $r > r_{\text{opt}}$ or $r \leq r_{\text{opt}}$. In the latter case, we must also give a set \mathcal{B} of k blue disk with radius r such that no blue disk overlaps any other disk. If we had an algorithm at our disposal that solves the decision problem in polynomial time, then we could solve the original optimization problem in polynomial time by applying Megiddo's parametric search [10]. Unfortunately, the decision problem is known to be \mathcal{NP} -complete [5]. Therefore we are looking for an algorithm that approximates the decision problem in the following sense: if m disks of radius r can be placed, then our algorithm places m disks of radius αr , for some fixed $\alpha \in (0, 1]$. If $m < k$, then we know that $r > r_{\text{opt}}$, and if $m \geq k$, then either $r \leq r_{\text{opt}}$, or $r > r_{\text{opt}}$ and $\alpha r \leq r_{\text{opt}}$. In other words, placement of at least k disks of radius less than αr_{opt} is guaranteed, and we may or may not be fortunate for radii in between αr_{opt} and r_{opt} .

Obviously, we would like to maximize α while keeping a polynomial running time. Given such an algorithm, we can use it to compute an α -approximation to the original optimization problem, again by using parametric search, albeit in a somewhat non-standard way.

By rescaling r to 1, we can regard the decision problem as that of packing $m \geq k$ unit disks into a rectangle that is already partially covered by n unit disks. In this paper, we consider the following problem:

Problem 1 (APPROXSIZESIZE) *Let $\alpha \in (0, 1]$ be a fixed real. Given a bounding rectangle P and a set \mathcal{R} of possibly intersecting unit disks whose centers lie in P , pack m non-intersecting disks of radius α into P , where m is the maximal number of unit disks that can be packed in P .*

Note that we do not know the value of m a priori. For $\alpha = 1/2$, the problem can be solved by placing disks with radius $1/2$ greedily, i.e., as long as there is space to place a disk, we place one at an arbitrary feasible position. The following simple charging argument shows that we will place at least m disk of radius $1/2$ in this way. Consider an arbitrary placement of m unit disks, and charge a disk C with radius $1/2$ to a unit disk D if the center c of C lies inside D . After the greedy algorithm has finished, all of the m unit disks have a charge of at least one. Otherwise, we can place a disk with radius $1/2$ in an uncharged unit disk such that their centers coincide, and this contradicts the termination condition of the greedy algorithm.

In their pioneering work [7] Hochbaum and Maas gave a polynomial-time approximation scheme (PTAS) for the problem of packing a maximal number

of unit disks into a region. The problem is known to be \mathcal{NP} -complete [5]. Even though the corresponding geometric dispersion problem looks very similar, inapproximability results have been shown. Baur and Fekete [2] proved hardness results for a variety of geometric dispersion problems, and their results can be modified to our setting with a bit of effort. Specifically, they showed that APPROXSIZE cannot be solved in polynomial time for any radius that exceed $13/14$, unless $\mathcal{P} = \mathcal{NP}$. Furthermore, for the case when the objects are squares, Baur and Fekete gave an $O(\log k \cdot n^{40})$ -time $2/3$ -approximation algorithm, where k is the number of squares. However, since a square is a simpler shape and easier to pack than a disk their approach cannot be generalized to disks. The main contribution of this paper is a polynomial-time $2/3$ -approximation algorithm. Actually, we conjecture that $2/3$ is indeed the largest value for which the problem can be solved in polynomial time.

APPROXSIZE has applications in non-photorealistic rendering system, where 3D models are to be rendered in an oil painting style, as well as in random examinations of, e.g., soil or water.

2 Algorithm outline

We now give a rough outline of our algorithm DISKPACKING. We use the term r -disk as shorthand for a disk of radius r . For $r > 0$ and a set $R \subseteq \mathbb{R}^2$ let the r -freespace $\mathcal{F}_r(R)$ of R be the set of the centers of all r -disks that are completely contained in R . By $\mathcal{F}_r^\otimes(R)$ we denote the Minkowski sum of $\mathcal{F}_r(R)$ and an r -disk.

We first compute the sets $\mathcal{F}_1 = \mathcal{F}_1(P \setminus \bigcup \mathcal{R})$ and $\mathcal{F}_1^\otimes = \mathcal{F}_1^\otimes(P \setminus \bigcup \mathcal{R})$, see Fig. 1. Then, we apply the PTAS of Hochbaum and Maas [7] to \mathcal{F}_1^\otimes . For any positive integer t , the PTAS packs in $O(n^{t^2})$ time at least $(1 - 1/t)^2 \cdot m$ unit disks into \mathcal{F}_1^\otimes , where m is the maximum number of unit disks that can be packed into \mathcal{F}_1^\otimes and n is the minimum number of unit squares whose union covers \mathcal{F}_1^\otimes . Setting $t = 25$ we obtain in $O(n^{625})$ time a set \mathcal{B} of $m' \geq 12/13 \cdot m$ unit disks.

Note that the approximation scheme by Hochbaum and Maas can be modified such that the algorithm is strongly polynomial with respect to the size of our input. If the number of disks that can be packed is not polynomial in the size of P and \mathcal{R} then there must exist a huge empty square region within P . This can be “cut out” and packed almost optimally by using a naïve approach. The added error obtained is bounded by $O(1/\tilde{n}^2)$ where \tilde{n} is the optimal number of disks that can be packed in the square. This step can be repeated until there are no more huge empty squares.

Given \mathcal{B} we compute a set $\mathcal{B}_{2/3}$ of disks of radius $2/3$ that has cardinality at least $13/12 \cdot m' \geq m$ and is contained in $P \setminus \bigcup \mathcal{R}$. We obtain $\mathcal{B}_{2/3}$ in two steps. First, we compute a sufficiently large matching in the nearest-neighbor graph $G = (\mathcal{B}, E)$ of \mathcal{B} with respect to a metric $\text{dist}(\cdot, \cdot)$ that we will specify later. Second, we define a region for each pair of matching unit disks such that we can place three $2/3$ -disks in each region (see Fig. 2) and all regions are pairwise disjoint. For each unmatched unit disk D we place a $2/3$ -disk D' such that the centers d and d' of D and D' , respectively, coincide.




Fig. 1. The 1-freespace \mathcal{F}_1^{\otimes} (light shaded) and a shortcut vv' (dashed) between the connected components C and C' of \mathcal{F}_1 .




Fig. 2. Packing three $2/3$ -disks in the region spanned by a pair of unit disks.

In the next sections we describe each step of Algorithm DISKPACKING in more detail.

3 The freespace and a metric on unit disks

We briefly recall the setting. We are given a set \mathcal{R} of unit disks whose centers lie in a rectangle P . The disks in \mathcal{R} are allowed to intersect. We first compute the freespace \mathcal{F}_1 of $P \setminus \bigcup \mathcal{R}$. According to Kedem et al. [8] the union of s disks can be computed in $O(s \log^2 s)$ time and its complexity is linear in s . Applying their algorithm to the disks in \mathcal{R} scaled by a factor of 2 and intersecting the resulting union with P , we can compute \mathcal{F}_1 in $O(|\mathcal{R}| \log^2 |\mathcal{R}|)$ time, where $|\mathcal{R}|$ is the cardinality of \mathcal{R} .

Next, we want to introduce a metric $\text{dist}(\cdot, \cdot)$ on unit disks in \mathcal{F}_1^{\otimes} . With the current definition of \mathcal{F}_1 we have the problem that two unit disks centered on points in different connected components of \mathcal{F}_1 can intersect. We solve this problem by considering a superset \mathcal{F}_1^+ of \mathcal{F}_1 that connects close components of \mathcal{F}_1 . By $|p, q|$ we denote the Euclidean distance between two points p and q in the plane.

Definition 1. Let \mathcal{C}_1 be the set of connected components of \mathcal{F}_1 , and let $C, C' \in \mathcal{C}_1$. Let v and v' be vertices on the boundaries of C and C' , respectively. We say that the line segment vv' is a shortcut if $|v, v'| \leq \sqrt{11} \cdot 2/3 \approx 2.21$. Let $\mathcal{S}(C, C')$ be the set of all shortcuts induced by C and C' . We set $\mathcal{F}_1^+ = \mathcal{F}_1 \cup \bigcup_{C, C' \in \mathcal{C}_1; s \in \mathcal{S}(C, C')} s$.

Figure 1 depicts \mathcal{F}_1 , \mathcal{F}_1^{\otimes} , and a shortcut vv' . Throughout the paper we will use upper-case letters to denote disks and the corresponding lower-case letters to denote their centers. Now, we are ready to define our metric for a connected component of \mathcal{F}_1^+ , see Fig. 3.

Definition 2. Let D and D' be unit disks in \mathcal{F}_1^\otimes , with centers d and d' , respectively. The distance $\text{dist}(D, D')$ of D and D' is the length of the geodesic $g(d, d')$ of d and d' in \mathcal{F}_1^+ . The tunnel $T(D, D')$ of D and D' is the union of all r -disks in $P \setminus \bigcup \mathcal{R}$ centered at points of $g(d, d')$ with $r \leq 1$.

It is easy to see that any $2/3$ -disk $D_{2/3}$ centered at a point of $g(d, d')$ does not intersect any disk in \mathcal{R} . (This will also follow from Lemma 2.) Thus $D_{2/3}$ is contained in the tunnel $T(D, D')$. The geodesic between two points in \mathcal{F}_1^+ can only consist of line segments and arcs of radius 2, see Fig. 3.

Recall that our algorithm computes a matching in the nearest-neighbor graph $G = (\mathcal{B}, E)$ induced by the metric $\text{dist}(\cdot, \cdot)$ on the set \mathcal{B} of unit disks that we get from the PTAS by Hochbaum and Maas. For each pair $\{D, D'\}$ in the matching we define a region $T_{2/3}(D, D')$ into which we will then place three $2/3$ -disks as in Fig. 2. An obvious way to define $T_{2/3}(D, D')$ would be to take the union of all $2/3$ -disks centered at points of the geodesic between d and d' in $\mathcal{F}_{2/3}$. Our definition is not as straight-forward, but will simplify the proof that $T_{2/3}(D, D')$ and $T_{2/3}(F, F')$ are disjoint if $D, D', F,$ and F' are pairwise disjoint. This is needed to ensure that the $2/3$ -disks that we will place in the tunnels $T_{2/3}$ are disjoint.




Fig. 3. The geodesic $g(d, d')$. Left: unrestricted. Center: obstacle R . Right: shortcut.

Definition 3. Let D and D' be unit disks in \mathcal{F}_1^\otimes . Let $g_{2/3}(d, d')$ be a geodesic from d to d' in $\mathcal{F}_{2/3}(T(D, D'))$. Then, the $2/3$ -tunnel $T_{2/3}(D, D')$ of D and D' is the union of all $2/3$ -disks centered at points of $g_{2/3}(d, d')$.

According to Chang et al. [4] the geodesics $g(d, d')$ and $g_{2/3}(d, d')$ from d to d' can be computed in $O(|\mathcal{R}|^2 \log |\mathcal{R}|)$ time.

4 The nearest-neighbor graph

Recall that m is the maximum number of disjoint unit disks that fit into \mathcal{F}_1^\otimes . For $t = 25$ the $(1 - 1/t)^2$ -approximation of Hochbaum and Maas [7] yields a set \mathcal{B} of $m' \geq 12m/13$ unit disks in \mathcal{F}_1^\otimes . Our plan is to compute the nearest-neighbor graph $G = (\mathcal{B}, E)$ induced by the metric $\text{dist}(\cdot, \cdot)$, find a matching of sufficient size in G , and finally place three $2/3$ -disks in the $2/3$ -tunnel $T_{2/3}(C, D)$ for each pair $\{C, D\}$ in the matching. We show that if we place another $2/3$ -disk for each unmatched disk in \mathcal{B} , then we place at least $13m'/12 \geq m$ disks of radius $2/3$ by this approach.

By construction, two unit disks D_1 and D_2 whose centers lie in different components of \mathcal{F}_1^+ have an empty intersection, so we can consider each connected component of \mathcal{F}_1^+ separately.

After running the algorithm of Hochbaum and Maas we greedily add to \mathcal{B} disjoint unit disks in $\mathcal{F}_1^\otimes \setminus \bigcup \mathcal{B}$ until no more disks can be added. This is needed to ensure the following lemma:

Lemma 1. *The nearest-neighbor graph $G = (\mathcal{B}, E)$ (w.r.t. dist) is planar and has maximum degree 6.*

Proof. Let $C \in \mathcal{B}$ be an arbitrary unit disk, let $C' \in \mathcal{B}$ be the nearest neighbor of C in \mathcal{B} , and let $\mathcal{D} = \{D_1, \dots, D_k\} \subseteq \mathcal{B}$ be the neighbors of C in \mathcal{B} for which C is their nearest neighbor. If $k \leq 5$ then the degree bound obviously holds, thus we only have to consider the case when $k \geq 6$. For each disk D_i , $1 \leq i \leq k$, place a unit disk D'_i with center on $g(c, d_i)$ such that $|c, d'_i| = 2$, i.e., D'_i touches C . From the definition of the nearest-neighbor graph it follows that every point on $g(c, d_i)$ is closer to C or D_i than to any other unit disk in $\mathcal{B} \setminus \{D_i\}$. As a result the set D'_1, \dots, D'_k and C' has to be disjoint. Using a simple packing argument, $k = 6$ and $C' \in \mathcal{D}$ follows and thus the degree bound stated in the lemma holds.

Finally, G is planar since no two edges in a nearest-neighbor graph can intersect. \square

From now on we will call $\{C, D\} \subseteq \mathcal{B}$ a *nearest pair* if either (C, D) or (D, C) is an edge in G , i.e., either D is the nearest disk in \mathcal{B} to C or C is the nearest disk in \mathcal{B} to D . For every nearest pair $\{C, D\}$ we define $\mathcal{A}(C, D)$ to be $C \cup D \cup T_{2/3}(C, D)$. As the nearest pair $\{C, D\}$ is a potential candidate to become a matching pair, we want to ensure that we can use $\mathcal{A}(C, D)$ to pack three $2/3$ -disks in it such that all the packed $2/3$ -disks are pairwise disjoint. Thus, we have to prove:

- (i) three $2/3$ -disks fit into $\mathcal{A}(C, D)$ and
- (ii) for any nearest pair $\{E, F\}$ where C, D, E and F are pairwise disjoint $\mathcal{A}(C, D) \cap \mathcal{A}(E, F) = \emptyset$.

Note that we do not have to care whether, e.g., $\mathcal{A}(C, D)$ intersects $\mathcal{A}(C, E)$ because the matching will choose at most one pair out of $\{C, D\}$ and $\{C, E\}$. Three $2/3$ -disks obviously fit into $\mathcal{A}(C, D)$ since C and D do not intersect, thus, (i) is fulfilled. The remaining part of the paper will focus on proving (ii).

We split the proof into two parts. The first part shows that $T_{2/3}(C, D)$ does not intersect any disk other than C and D . The second part shows that no two $2/3$ -tunnels $T_{2/3}(C, D)$ and $T_{2/3}(E, F)$ intersect. We start with two technical lemmas that we need to prove the first part.

Lemma 2. *Let C and D be two unit disks in \mathcal{F}_1^\otimes . If $|c, d| \leq \frac{2}{3}\sqrt{11}$ then $g_{2/3}(c, d)$ is a line segment.*

Proof. Let $T'_{2/3}(C, D)$ be the Minkowski sum of a $2/3$ -disk and the line segment cd , see Fig. 4a. If $g_{2/3}(c, d)$ is not a line segment, then a disk E in $\mathcal{B} \cup \mathcal{R}$ intersects $T'_{2/3}(C, D)$. We establish a lower bound on $|c, d|$ for this to happen. Note that C, D and E are pairwise disjoint as C and D are disks in \mathcal{B} .

Clearly, the minimum distance between c and d is attained if E and $T'_{2/3}(C, D)$ only intersect in a single point and furthermore, both E and C as well as E and D intersect in a single point. This means that $|c, e| = |d, e| = 2$. Moreover, the Euclidean distance between e and the straight-line segment cd is $1 + \frac{2}{3} = \frac{5}{3}$. By Pythagoras' theorem we calculate $|c, d|$ to be at least $\frac{2}{3}\sqrt{11}$. This means that $T'_{2/3}(C, D)$ is contained in $P \setminus \mathcal{R}$. Even if C and D belong to different components of \mathcal{F}_1 , by Definition 1 they are connected via a shortcut. Thus, $g_{2/3}(c, d)$ is a line segment. \square

Lemma 3. *Let D and E be two unit disks in \mathcal{F}_1^\otimes that are infinitesimally close to each other. Then $\text{dist}(D, E) \leq \frac{2}{3}\pi$.*

Proof. For simplification we assume that D and E touch, as illustrated in Fig. 4a. The curve $g(D, E)$ attains its longest length if there is an obstacle disk R that touches D and E and no shortcut could be taken. In this case $g(D, E)$ describes an arc of radius 2 and 60° , thus its length is $\frac{1}{6} \cdot 2 \cdot 2\pi = \frac{2}{3}\pi$. \square

Now, we are ready to prove the first part:

Lemma 4. *Let $\{C, D\} \subseteq \mathcal{B}$ be a nearest pair. No disk of $\mathcal{B} \cup \mathcal{R} \setminus \{C, D\}$ intersects $T_{2/3}(C, D)$.*

Proof. From the definition of freespace and Definitions 2 and 3 it immediately follows that neither $T(C, D)$ nor $T_{2/3}(C, D)$ are intersected by a disk in \mathcal{R} . Thus, it remains to prove that apart from C and D no disk in \mathcal{B} intersects $T_{2/3}(C, D)$.

W.l.o.g. let C be the nearest disk in \mathcal{R} to D . The proof is done by contradiction: we assume that there is a disk $E \in \mathcal{B}$ that intersects $T_{2/3}(C, D)$.

First, we move a unit disk on $g(C, D)$ from the position of D to the first position in which it hits E , denote the disk in this position by \bar{D} , see Fig. 4a where $D = \bar{D}$ holds. Note that \bar{D} does not necessarily lie entirely within \mathcal{F}_1^\otimes . However, according to Lemma 2 (C, \bar{D} and E are disjoint), the Euclidean distance between c and \bar{d} is at least $\frac{2}{3}\sqrt{11}$. We prove that the geodesic $g(\bar{d}, e)$ within \mathcal{F}_1^+ is of length less than $g(c, \bar{d})$. This contradicts C being the nearest neighbor of D .

In the remainder of the proof we show that $g(c, \bar{d})$ is larger than $g(d, \bar{e})$, regardless of whether \bar{d} is in \mathcal{F}_1 or not. The details, especially of the latter case, are rather technical and somewhat unpleasant, and due to space limitations we omit them here. They can be found in the full version of this paper [1]. \square

Lemma 4 settles that no other disks apart from C and D intersect $T_{2/3}(C, D)$. We still have to show that no two $\frac{2}{3}$ -tunnels $T_{2/3}(C, D)$ and $T_{2/3}(E, F)$ intersect.

Theorem 1. *Let $\{C, D\}, \{E, F\} \subseteq \mathcal{B}$ be two nearest pairs such that C, D, E and F are pairwise disjoint, it holds that $T_{2/3}(C, D) \cap T_{2/3}(E, F) = \emptyset$.*

Proof. The proof is by contradiction again. Assume that $T_{2/3}(C, D)$ and $T_{2/3}(E, F)$ intersect. First, we exclude the scenario in which even the geodesics $g_{2/3}(c, d)$ and $g_{2/3}(e, f)$ intersect. If $g_{2/3}(c, d)$ and $g_{2/3}(e, f)$ intersect it immediately follows that $g(c, d)$ and $g(e, f)$ also intersect. Note that this comprises the case in which $g(c, d)$ and $g(e, f)$ take the same shortcut. Let i be one of the intersection points of $g(c, d)$ and $g(e, f)$. By appropriately exchanging parts of $g(c, d)$ and $g(e, f)$ having endpoints c, d, e, f and i , it is easy to construct a curve that at least contradicts one of $g(c, d)$ and $g(e, f)$ to be a geodesic and thus one of the pairs $\{C, D\}$ or $\{E, F\}$ to be nearest. Thus, we can assume that only $T_{2/3}(C, D)$ and $T_{2/3}(E, F)$ intersect. Obviously, it is enough to prove the theorem for the case in which the tunnels intersect in a single point p , see Fig. 4b. Thus, p lies in \mathcal{F}_1 and neither $g(c, d)$ nor $g(e, f)$ takes a shortcut containing p and we can w.l.o.g. assume that no shortcut is taken at all (comparing with one of the disks of $\mathcal{D}(S)$ on $g(c, d)$ or $g(e, f)$ instead of C, D, E or F for the used shortcut S). Again, we show that $\{C, D\}$ and $\{E, F\}$ can't be nearest pairs at the same time.

We observe that at least one of the disks $\{C, D, E, F\}$ intersects the unit disk P with center p ; otherwise there would be another disk in \mathcal{B} located in the space between C, D, E and F which would immediately contradict $\{C, D\}$ as well as $\{E, F\}$ being nearest pairs. W.l.o.g. let C be a disk that intersects P .

Let p_{CD} be the point on $g_{2/3}(C, D)$ such that $|p, p_{CD}| = 2/3$, see Fig. 4b. Define p_{EF} correspondingly. We assume that there is a vicinity of p_{CD} and p_{EF} in which $g_{2/3}(C, D)$ and $g_{2/3}(E, F)$ are arcs. The case when one vicinity of p_{CD} and p_{EF} is a straight-line is easier and can be treated with similar arguments.

The curvature of $g_{2/3}(C, D)$ and $g_{2/3}(E, F)$ in a vicinity of p_{CD} and p_{EF} induces the existence of two disks $R, S \in \mathcal{R}$ as illustrated in Fig. 4b. Since R and S forces the curvature of $g_{2/3}(C, D)$ and $g_{2/3}(E, F)$ we may introduce the following coordinate system. The origin is p and the coordinates of r and s are $(0, \frac{7}{3})$ and $(0, -\frac{7}{3})$, respectively.

As a consequence of Lemma 2 we get that each geodesic $g_{2/3}$ starts with a straight-line segment of length at least $\frac{1}{3}\sqrt{11} \approx 1.11$. Thus, the curvature of $g_{2/3}(C, D)$ in p_{CD} infers that $|c, p_{CD}| \geq 1.11$ holds, which means that C either lies completely to the left of the y -axis or to the right. This holds analogously for the other disks. W.l.o.g. we assume that C and E lie to the left of the y -axis and D and F lie to the right, see Fig. 4b.

Note that we have to take care which relationship inferred that $\{C, D\}$ and $\{E, F\}$ are nearest pairs, e.g. C could be the nearest neighbor of D or D could be the nearest neighbor of C . We will prove the following:

- (i) $\text{dist}(C, E) < \text{dist}(E, F)$
- (ii) $\text{dist}(C, E) < \text{dist}(C, D)$
- (iii) $\text{dist}(D, F) < \text{dist}(C, D)$

Item (i) says that C is closer to E than F is. Thus, in order for $\{E, F\}$ to be a nearest pair, E must be the nearest neighbor of F . We use this fact to show that (ii) and (iii) hold. Together, (ii) and (iii) comprise the contradiction: (ii) says that D is not the nearest neighbor of C , while (iii) says that C is not the nearest neighbor of D . Hence, $\{C, D\}$ cannot be a nearest pair.

The proofs of (i)—(iii) can be found in the full version of this paper [1]. \square




Fig. 4. Illustrations for (a) the proof of Lemma 2 and 3. (b) the proof of Theorem 1.

5 The set $\mathcal{B}_{2/3}$

After computing \mathcal{B} and the nearest neighbor graph $G = (\mathcal{B}, E)$, we compute a matching in G . Let $m' = |\mathcal{B}|$ be the number of unit disks in \mathcal{B} . Recall that G is planar and has degree at most 6. We show that we can find a matching in which the number of matched disks is at least $\frac{1}{6} \cdot m'$. Observe that G can consist of more than one connected component. We look at each connected component separately. Let C be a connected component and let c be the number of disks that it contains. Clearly, C contains a spanning tree of degree at most 6. It is easy to see that there is a matching in C that matches at least $\frac{1}{6} \cdot c$ disks. Doing this for each connected component yields a matching in G that contains at least $\frac{1}{6} \cdot m'$ matched disks.

According to Theorem 1 and Lemma 4 we can pack three $2/3$ -disks in $\mathcal{A}(C, D)$ for every matched pair $\{C, D\}$ such that these $2/3$ -disks are pairwise disjoint. For each of the remaining unmatched disks we pack one $2/3$ -disk in each disk. Let set of all disks packed as above be $\mathcal{B}_{2/3}$. By construction, there are no interferences between these sets belonging to different connected components of

\mathcal{F}_1^+ . The cardinality of $\mathcal{B}_{2/3}$ is at least $\frac{1}{6} \cdot \frac{3}{2} \cdot m' + \frac{5}{6} \cdot m' = \frac{13}{12} \cdot m'$. Since the cardinality of \mathcal{B} is at least $\frac{12}{13} \cdot m$, the set $\mathcal{B}_{2/3}$ contains at least m $2/3$ -disks and we can conclude with the following theorem:

Theorem 2. *Algorithm DISKPACKING is a polynomial-time $2/3$ -approximation for the problem APPROXSIZE.*

6 Conclusion

Naturally, our result is purely of theoretic interest. The bottleneck for the running time is the application of Hochbaum and Maas' PTAS with approximation factor $(1 - 1/13)$. To obtain an algorithm with better running time, it seems unavoidable to use a completely different approach. For future work it would also be desirable to narrow the gap between the approximation factor of $2/3$ of our algorithm and the inapproximability result of $13/14$ of Baur and Fekete [2]. We conjecture that, unless $\mathcal{P} = \mathcal{NP}$, the lower bound of $2/3$ is indeed tight.

References

1. M. BENKERT, J. GUDMUNDSSON, C. KNAUER, E. MOET, R. VAN OOSTRUM AND A. WOLFF. A polynomial-time approximation algorithm for a geometric dispersion problem. Technical Report 2005-8, Universität Karlsruhe, May 2006. Available at <http://www.ubka.uni-karlsruhe.de/indexer-vvv/ira/2006/8>
2. C. BAUR AND S. P. FEKETE. Approximation of Geometric Dispersion Problems. *Algorithmica*, 30:450–470, 2001.
3. P. CAPPANERA. A survey on obnoxious facility location problems. Technical Report TR-99-11, University of Pisa, 1999.
4. E.-C. CHANG, S. W. CHOI, D. KWON, H. PARK AND C.-K. YAP. Shortest path amidst disc obstacles is computable. In proceedings of the 21st *ACM Symposium on Computational Geometry*, 2005.
5. R. J. FOWLER, M. S. PATERSON AND S. L. TANIMOTO. Optimal packing and covering in the plane are NP-compet. *Information Processing Letters*, 12:133–137, 1981.
6. Z. FÜREDI. The densest packing of equal circles into a parallel strip. *Discrete & Computational Geometry*, 6:95–106, 1991.
7. D. HOCHBAUM AND W. MAAS. Approximation Schemes for Covering and Packing Problems in Image Processing and VLSI. *Journal of the ACM*, 32:130–136, 1985.
8. K. KEDEM, R. LIVNE, J. PACH AND M. SHARIR. On the union of Jordan regions and collision-free translational motion amidst polygonal obstacles. *Discrete & Computational Geometry*, 1:59–71, 1986.
9. C. MARANAS, C. FLOUDAS AND P. PARDALOS. New results in the packing of equal circles in a square. *Discrete Mathematics*, 128:187–293, 1995.
10. N. MEGIDDO. Applying parallel computation algorithms in the design of serial algorithms. *Journal of the ACM*, 30(4):852–865, 1983.
11. C. A. ROGERS. Packing and Covering. *Cambridge University Press*, 1964.
12. G. F. TÓTH. Packing and Covering. In Handbook of Discrete and Computational Geometry, 2nd Edition J. E. Goodman and J. O'Rourke, editors CRC Press LLC, 2004.
13. C. ZONG AND J. TALBOT. Sphere Packings. Springer-Verlag, 1999.